

Automatic Detection and Classification of Boundary Crossings in Spacecraft in situ Data

H.T. Rüdissler^{1,2}, A. Windisch^{1,4}, U.V. Amerstorfer³,
D. Píša⁵, J. Soucek⁵

¹Know Center, Graz, Austria; ²University of Graz, Graz, Austria;

³Space Research Institute, Austrian Academy of Sciences, Graz, Austria;


⁴Department of Physics, Washington University in St. Louis, USA;

⁵Institute of Atmospheric Physics of Czech Academy of Sciences, Prague, Czechia

Planetary magnetospheres create multiple sharp boundaries, such as the bow shock, where the solar wind plasma is decelerated and compressed, or the magnetopause, a transition between solar wind field and planetary field. We attempt to use convolutional neural networks (CNNs) as well as traditional machine learning algorithms to identify magnetospheric boundaries, i.e. planetary and interplanetary shocks crossings and magnetopause crossings in spacecraft in situ data. The boundaries are identified by a discontinuity in a magnetic field, plasma density, and in the spectrum of high-frequency waves. These measurements are available on many planetary missions. Data from Earth's missions Cluster and THEMIS are used for CNN training. We ultimately strive for successful classification of boundaries (shock, magnetopause, inbound, outbound) and the correct handling of multiple crossings.

Planetary magnetospheres create multiple sharp boundaries

- ▶ Bow shock, where the solar wind plasma is decelerated and compressed
- ▶ Magnetopause – a transition between solar wind field and planetary field



Solar wind and magnetosphere. Image: ESA


Shocks can be identified in multiple datasets:

- ▶ Magnetic field
- ▶ Ion density and speed
- ▶ Ion spectrum
- ▶ Electric field spectrum

Signatures of a shock:

- ▶ B increases
- ▶ Velocity decreases
- ▶ Density increases
- ▶ Electric field spectrum and ion spectrum change character

Example of a bow shock crossing. Data from WHISPER C1 and CIS-HIA-HS C1.



There exist multiple spacecraft providing datasets for the identification of bow shock crossings.

- ▶ 4 Cluster spacecraft (since 2001, will operate atleast until the end of 2022)
- ▶ 5 THEMIS spacecraft (operational since 2007)

Usually, the shock is crossed by all 4 Cluster spacecraft. The relative timing of these crossings is valuable information for 3D modelling, but human expert labeling is time consuming. Since not all shocks look exactly the same and especially quasi parallel shocks can exhibit less sharp signatures, threshold based algorithms are not sufficient to reliably detect shocks.

Convolutional neural networks (CNNs) are commonly used for computer vision tasks, such as object detection. Nevertheless, patterns in time series can be detected as well. Large amounts of data can be processed in short time, once the training is completed.

Steps so far:

- ▶ Preprocessing Cluster data for training
- ▶ Labeling and defining NN architecture

Next steps:

- ▶ Training on Cluster Data
- ▶ Hyperparameter tuning
- ▶ Testing algorithm on new, unseen data

Ultimately:

- ▶ Expanding possibility of algorithm to detect and label different regions
- ▶ Testing other machine learning methods on this use case

Europlanet 2024 RI has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 871149.

A. Windisch acknowledges support by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under Award Number # DE-FG-02-05ER41375.